Index to Volume 89 of THE BRYOLOGIST[1,2]

4-*O*-methylolivetoric acid, 297
acetyl-polymalonyl derived secondary products in lichens, 92
acetyl-polymalonyl pathway, in lichen systematics, 91+
Acrolejeunea, 186
Acrobolbus ochrophyllus, 294
Acroporium, 26; pungens, 26; sehnemii, 10; **ulicinum**, comb. nov., 26
Actinodontium portoricense, 11
Actinogyra muhlenbergii, *285*
Adelanthus lindbergianus, 293; tenuis, 293
Africa, elements in Venezuela, 11; first report of Syrrhopodon incompletus in, 213
AGARWAL, A. *See* SRIVASTAVA, S. C. & A. AGARWAL, 195
AHMADJIAN, V. (review). G. Schuster, Die Jugendentwicklung von Flechten, 185
Alectoria, 102
aliphatic compounds, 91
Allantoparmelia, 219+
allelopathy, lichen substances in, 113
ALLEN, B. H. (review). J.-P. Frahm, Africanische Campylopus-Arten (Dicranaceae, Musci), 248
ALLEN, B. H. (review). S. P. Churchill, A Synopsis of the Kansas Mosses with Keys and Distribution Maps, 185
ALLEN, B. H. A review of the genus Parisia (Musci: Dicranaceae), 206
ALLEN, B. H. The taxonomic status of Hypnella punctata, 224
Allisoniella, 155; subbipartita, 293+
Amblystegium serpens, 61
Amphicephaloziella, 155
Amygdalaria, 136
Anaptychia, 102; fusca, 117
Anastrophyllum ciliatum, 294; involutifolium, 293; schismoides, 293
Andes, Venezuelan bryoflora of, 8
ANDO, H. *See* NISHIMURA, N. & H. ANDO, 66
Andreaea alpestris, 193; alpina, 193; angustata, *192*; blyttii, 192+; blyttii ssp. angustata, *192*; heinemannii, 192+; key to ecostate species of North America and Europe, 193; marginata, 194; mutabilis, 193; obovata, 193; rupestris var. papillosa, 193; rupestris var. rupestris, 189+; **sinuosa**, sp. nov., 189+, 190*
Anisothecium campylophyllum, 10
Anoectangium obtusifolium, 33
Anomobryum plicatum, 9
Anomodon attenuatus, 60+; minor, 60+; rostratus, 60+
antibiotics, lichen substances as, 112
antiherbivory, lichen secondary products in, 115
Aongstroemia julacea, 9
Aptychella proligera, 8
Archidium, 42+, 144+; tenerrimum, 144+; blepharoplast morphology in, 42, 44*, 45*, 47*, 48*
Arcto-Tertiary Geoflora, refugium in Interior Highlands, 32
aromatic compounds, 91
Arthonia palmulacea, 232+
Asia, elements in the Interior Highlands bryoflora, 32
Aspicilia cinerea, 116; gibbosa, 116
Astomum ludovicianum, 33
Astroplaca, 102
Atractylocarpus costaricensis, 33
Atrichum crispum, 33
Aulaxina quadrangula, 233

Baeomyces rubiformis, *271*
Balantiopsis asymmetrica, 294
Barbula ehrenbergii, 33; indica, 33; replicata, 10; spiralis, 10; unguiculata, 70, 227
Bartramia angustifolia, 9
BASILE, D. V. Dedication, Aaron J. Sharp Dedication Issue, 1
BAUER, C. R. *See* CROSBY, M. R. & C. R. BAUER, 168
Bazzania trilobata, 32
Biatora crenata, 253; decipiens, 253; globifera, 253; globifera var. rubiformis, 253; himalayana, *262*; icterica, *253*+; luridella, 253; russellii, 253
biogenetic hypotheses in lichen systematics, 91
biography, A. J. Sharp, 3
Blepharidophyllum clandestinum, 292+; densifolium, 292
blepharoplast morphology, in Archidium tenerrimum, 42; in Funaria hygrometrica, 46; in Thuidium delicatulum, 144+
bourgeanic acid, 93
Brachelyma subulatum, 33
Brachydontium, 220+; flexisetum, 200+; intermedium, 200+; **olympicum**, comb. nov., 200; trichodes, 200+
Brachythecium acuminatum, 61+; oxycladon, 61
Braunfelsia, 206+; enervis, 206
Brazil, Fissidens austro-americanus, F. brittonii, F. neglectus, F. reesei new to, 300
Breedlovea chiapensis, gen. et sp. nov., 23, 24*
Breutelia allionii, 8
BRODO, I. M. & W. L. CULBERSON. Haematomma pustulatum, sp. nov. (Ascomycotina, Haematommataceae): a common, widespread, sterile lichen of eastern North America, 203
BRODO, I. M. Interpreting chemical variation in lichens for systematic purposes, 132
Brodoa, gen. nov., 219+; **atrofusca**, comb. nov., 222; **intestiniformis**, comb. nov., 222; **oroarctica**, comb. nov., 222
Brothera leana, 32
Brotherella tenuirostris, 32
Brotherobryum, 206+
BROWN, C. R. & Z. B. CAROTHERS. Comparative studies of spermatogenesis in the Bryopsida. II. Blepharoplast morphology in Archidium tenerrimum Mitt., 42
Bruchia hallii, 33
Bryoandersonia illecebra, 60
Bryoerythrophyllum, 13; andersonianum, 15; arcuatum, *15*; binnsii, 15; bolivianum, 14; calcareum, 13+; caledonicum, 16; campylocarpum, 14; ferruginascens, 14; inaequalifolium, 14; jamesonii, 9, 14; key to Mexican species, 15; lusitanicum, *15*; recurvirostrum, 15; recurvirostrum var. robustum, 16; recurvum, 14; **sharpii**, sp. nov., 13*; wallichii, 16
Bryoria, 102
Bryoxiphium norvegicum ssp. norvegicum, 33
Bryum andicola, 227; **calobryoides**, sp. nov., 215, 216*; capillare, 215; capitellatum, *215*; chryseum, 9; cyclophyllum, 33, 217; gerwigii, 217; incrassatolimbatum, 33; miniatum, 33; viridulum, 36

---

[1] Compiled by P. M. Eckel, Clinton Herbarium, Buffalo Museum of Science, Buffalo, NY 14211.

[2] New scientific names and new combinations are printed in boldface type; illustrations are indicated by an asterisk; taxa discussed or mentioned several times in a paper are indicated by a plus sign; italicized page numbers indicate taxa in synonymy.

BUCK, W. R. (review). H. C. Gangulee, Handbook of Indian Mosses, 250
BUCK, W. R. (review). L. P. Thomas & J. R. Jackson, Walk Softly Upon the Earth. A Pictorial Field Guide to Missouri Mosses, Liverworts and Lichens, 249
Bulbothrix, 102
Byssoloma subdiscordans, 233; tricholomum, 233

Calenia aspidotum, 232
Callicladium haldanianum, 32
Callicosta, 11
Caloplaca citrina, 115
Calymperes, 49; motleyi, 50
Calypogeja muelleriana subsp. blomquistii, 33; sullivantii, 32
CAMERON, R. G. & R. WYATT. Substrate restriction in entomophilous Splachnaceae: role of spore dispersal, 279
Campylium hispidulum, 61
Campylopidulum, 152
Campylopodium, 23, 152; pussillum, 8
Campylopus, 152; annotinus, 8; apollinairei, 9; basalticolus, *153*; bogotensis, 153; cucullatifolius, 8; hoffmanii, 11; pilifera, 33; saxatilis, 9; strictulus, 8; subchlorophyllosus, *152*+; talluensis, 33
Campylostelium saxicola, 33
Cape Horn, hepatics reported for, 291+
CAROTHERS, Z. B. *See* BROWN, C. R. & Z. B CAROTHERS, 42
CAROTHERS, Z. B. *See* RUSHING, A. E. & Z. B. CAROTHERS, 144
Catillaria bouteillei, 233
Cavernularia, 219+
Central America, Elmeriobryum guatemalense in, 28
Cephalojonesia, 155
Cephalozia connivens, 60
Cephaloziella, 155+; alpina, *155*; arctica, 155+; brinkmanii, 155+; byssaceae, 155; dentata, 161; divaricata, 155+; divaricata var. scabra, 155+; elachista, 155+; hampeana, 155+; key to western North American species, 155; myriantha, *155*; papillosa, *155*; papillosa var. sinuata, *155*; patulifolia, 155+; phyllacantha, 156; rubella, 155+; stellulifera, 155+; stellulifera var. gracillima, 155+; striatula, *155*; subdentata, 155+; turneri, 155+
Cephaloziellaceae, in western North America, 155+
Cephaloziopsis, 155
Cetraria arenaria, 139; ciliaris, 104; ericetorum, 108; halei, 104; islandica, 107+, 117; microphyllica, 104; nivalis, 117; orbata, 104
Cetrariastrum, 129; sorocheilum, 129
Cetrelia, 102; evolution of secondary products in, 93
Cheilolejeunea, 195; warnstorfii, 292+
Cheilothela vaginata, 10
chemosyndrome variation, in Xanthoparmelia species, 100
chemotypes, as sibling species in lichens, 126
Chiloscyphus gottscheoides, 292; hookeri, 292+; pallidovirens, 292, 293; porrectus, 292, *293*
Chlorisodontium wallisii, 8
Chryso-hypnum diminutivum, 66, 67*; elegantulum, 66+; pendulinum, *69*; **squarrosulum**, comb. nov., 67*+
Cladia aggregata, 108; ferdinandii, 107
Cladina evansii, 100; impexa, 117; subtenuis, 119
cladogram, chemotypes of Cladonia chlorophaea group, 96
Cladomastigum bifidum, 11
Cladonia, 134; asahinae, 95; cariosa, 104; chlorophaea, 97; chlorophaea group, 91+, 105, chemical analysis of, 91, cladogram of chemotypes of, 96, siblings species in, 126; conista, 95+; cristatella, 115, 139+, 285, enzyme analysis of morphotypes in, 141; cryptochlorophaea, 96; fimbriata, 97; foliacea, 117; grayi, 96; homosekikaica, 96; imbricarica, 96; multiformis, 100; novochlorophaea, 96; perlomera, 96; polycarpia, 104, 128; polycarpoides, 104, 128; subconistea, 95
Clasmatocolea fulvella, 293; humilis, *292*+; obvoluta, 292+; vermicularis, 292
Clasmatodon parvulus, 60+
Cnestrum, 24
Coastal Plain, elements in the Interior Highlands, 33
Coccotrema pocillarium, 136
Collemataceae, 100
Cololejeunea biddlecomiae, 32
Colombia, Fissidens neglectus new to, 300
communities, corticolous bryophyte, 59
Conardia compacta, 32
Cornicularia aculeata, 117
Coscinodon arsenei, *24*
CRANDALL-STOTLER, B. *See* STOTLER, R. E. & B. CRANDALL-STOTLER, 73, 168
Crassicosta hermannii, 227
CROSBY, M. R. & C. R. BAUER. Recent literature on mosses—135, 177
CRUM, H. Taxonomic and nomenclatural addenda to the Mexican moss flora, 23
Cryphaea glomerata, 33; ravenelii, 33
Cryptochila grandifolia, 292
CULBERSON, C. F. Biogenetic relationships of the lichen substances in the framework of systematics, 91
CULBERSON, W. L. Introduction. Natural-product chemistry in the systematics of the lichen fungi, 89
CULBERSON, W. L. (review). H. Hertel & F. Oberwinkler (eds.). Beiträge zur Lichenologie. Festschrift J. Poelt, 86
CULBERSON, W. L. (review). P. F. Cannon, D. L. Hawksworth & M. A. Sherwood-Pike, The British Ascomycotina, an Annotated Checklist, 86
CULBERSON, W. L. Chemistry and sibling speciation in the lichen-forming fungi: ecological and biological considerations, 123
CULBERSON, W. L. *See* BRODO, I. M. & W. L. CULBERSON, 203
Cygniella sharpii, gen. et sp. nov., 23, 24*
Cylindrocolea, 155

DELGADILLO M., C. Aaron J. Sharp: a life-time of botanical experience, 3
depsidones, origin in lichens of, 93
Desmatodon plinthobius, 33
Dichelyma capillaceum, 32
Dicnemonaceae, 206
Dicranella cardotii, 10
Dicranoloma, 207
Dicranum basalticolum, *153*; montanum, 61+; polysetum, 33; scoparium, 61
Didymodon acutus var. acutus, 32; australasiae var. australasiae, 227; australasiae var. umbrosus, 70+, 71*; calymperidictyon, *15*; campylocarpus, *15*; linearis, *15*; revolutus, 72; rigidulus, 32; rigidulus var. gracilis, 227; rigidulus var. rigidulus, 70+
Dimelaena oreina, 105
Diplophyllum apiculatum var. apiculatum, 33
disjunction, Andreaea sinuosa, western North America to western Europe, 191; Syrrhopodon incompletus var. incompletus. Americas to Africa, 213; Entosthodon laxus, circum-subantarctic to Andes, 308

ECKEL, P. M. Didymodon australasiae var. umbrosus new to eastern North America, 70
Ecuador, bryoflora of, 17; Fissidens ramicola new to, 300; Lindigia debilis in, 230; Trachyodontium zanderi in, 17
EGAN, R. S. Correlations and non-correlations of chemical variation patterns with lichen morphology and geography, 99
EGAN, R. S. Recent literature on lichens—126, 80; —127, 234; —128, 310
elatinic acid, 203
Elharveya, nom. nov., 26; **vesicularioides**, comb. nov., 26
Elmeriobryum, 30; **guatemalense**, sp. nov., 29, 30*; philippinense, 30
Encalypta procera, 33
endemism, Appalachians and Interior Highlands, 32; West Indies, 11
Endocarpon crenatum, *260*
Entodon macropodus, 33; seductrix, 61
Entosthodon, 305; acutifolius, 305; antarcticus, *303*+; attenuatus, 304; bonplandii, 304; fascicularis, 305; fontanus, *303*+; gracilis, 305; laxus, 302+, 306*; longicollis, 305; longisetus, *305*; obtusifolius, 305; subattenuatus, *304*; vacifolius, *304*; venezuelanus, *303*+; wichurae, 305
Eopleurozia paradoxa, 9
Eriodon brevisetus, *230*
Erythrophyllopsis, 13; andina, 14
ESSLINGER, T. L. Notes on the chemistry and distribution of selected Parmeliaceae (lichens) from the southern hemisphere, 296
Eucamptodon ciliatus, *206*; laevipilus, *206*; macrocalyx, *206*
Eucamptodontopsis pilifera, 11
Eucladium verticillatum, 33
Eulejeunea olivacea, *195*
Eurhynchium mexicanum, *69*
Evernia prunastri, 117+

Fabronia ciliaris, 32
FAHSELT, D. Multiple enzyme forms of morphotypes in a population of Cladonia cristatella Tuck., 139
FAHSELT, D. *See* HAGEMAN, C. & D. FAHSELT, 285
fatty acids, 91+; in the evolution of orcinol compounds, 95
FIFE, A. J. Taxonomic and nomenclatural observations on the Funariaceae. 4. A review of Entosthodon laxus with incidental notes on E. obtusifolius, 302
Fissidens abruptus, *300*+; adianthoides, 36; asplenioides, 36+; austro-americanus, 300; bourgaeanus, 38; brevipes, 300; brittonii, 300; bryoides, 36+; cristatus, 60; exilis, 36+; gardneri, 300; gymnogynus, 37; incompletus, 39, 252; intermedius, 301; macroblastus, 300; micropyxis, 301; neglectus, 301; nitidulus, 22; osmundoides, 36+; palmatus, 36+; polypodioides, 36+; pseudomicrocladus, 301; ramicola, 300+; reesei, 301; semicompletus, 36+; serrulatus, 38; subbasilaris, 36+; taxifolius, 36+; typification of Hedwig's species of, 35; weirii, 10; zollingeri, 37
Flavopunctelia, 102+
Fontinalis sphagnifolia, 32
Forsstroemia ohioensis, 32; trichomitria, 32, 60
FRAHM, J.-P. A review of Sphaerothecium Hampe, 152
FRISVOLL, A. A. The status of Schistidium andreaeopsis, with notes on the Krause collection from the Chuck Peninsula, 276
Frullania eboracensis, 60; fertilis, 295; nisquallensis, 291
Funaria antarctica, 303; cavifolia, *303*+; hygrometrica, 114+, 144, 227+, blepharoplast morphology in, 46+; laxa, *304*; producta, 303; pungens, *303*+; subattenuata, *303*+; subcuspidata, 304

Gackstroemia decipiens, 294; magellanica, 294
galactinulin, 136
Geocalyx graveolens, 33
GLIME, J. M. *See* RAEYMAEKERS, G. & J. M. GLIME, 318
Glossadelphus, 11
Gollania, 30
GOWARD, T. Brodoa, a new lichen genus in the Parmeliaceae, 219
GRADSTEIN, S. R. Correction to review of R. Grolle's Nomina Generica Hepaticarium, 186
Graphidaceae, 100
Graphina confluens, 107
Graphis scripta, 115
GRIFFIN III, D. Rhynchostegiella attenuata Bartr. (Brachytheciaceae): a synonym of Lindigia aciculata (Tayl.) Hampe (Meteoriaceae), 230
GRIFFIN, D. *See* LEWINSKY, J. & D. GRIFFIN, III, 163
Grimmia alpestris, 200; andraeopsis, *276*+; arsenei, 24; **churchilliana**, nom. nov., 24; olneyi, 32; olympica, *200*+; pulvinata, 32; raui, 33; teretinervis, 33; wrightii, 33
Guayana Highlands, bryoflora of, 10
Guembelia, 200+
Gyalectidium filicinum, 233
Gymnostomum aeruginosum, 227; laxum, *302*+
Gyroweisia, 227+; **monterreia**, sp. nov., 227, 228*; reflexa, 228

HÄSSEL DE MENÉNDEZ, G. G. The discovery of Sullivant's Wilkes U.S. Exploring Expedition hepaticae. The "Fuegian" specimens, 291
Haematomma elatinum, 203+; lapponicum, 104; ochrophaeum, 203; **pustulatum**, sp. nov., 203, 204*; ventosum, 104
HAGEMAN, C. & D. FAHSELT. A comparison of isozyme patterns of morphological variants in the lichen Umbilicaria muhlenbergii (Ach.) Tuck., 285
Haplohymenium triste, 61+
Haplomitrium, 47
Harveya, *26*; vesicularioides, *26*
HEDDERSON, T. A. A naturally occurring moss hybrid between Orthotrichum gymnostomum and O. obtusifolium from Newfoundland, Canada, 165
Hedwig-Schwaegrichen Herbarium, 35
Helodium paludosum, 32
HERMANN, F. J. *See* ZANDER, R. H. & F. J. HERMANN, 227
Herpetineuron toccoae, 33
Herzogiella striatella, 33
Heterodermia, 102
Heteroscyphus integrifolius, 292
Holomitriopsis laevifolia, 11
Holomitrium flexuosum, 9+; marginatum, 10
Homalia defoliata, 10; trichomanoides, 32
Homaliadelphus sharpii, 33
Homalotheciella subcapillata, 33
HONG, W. S. The family Cepaloziellaceae in North America west of the hundredth meridian, 155
Hookeria acutifolia, 33; leptorrhyncha, *224*; punctata, *224*
Hookeriopsis, 10; **bowersiana**, sp. nov., 25*; cruegeriana, 11, 25; diffusa, 25; guadalupensis, 10, 26; heteroica, 25*+; maguirei, 11; obtusifolia, 9; **welchiana**, sp. nov., 25*
Husnotiella revoluta var. palmeri, 33
hybrid, Orthotrichum gymnostomum and O. obtusifolium, 165, 166*
Hymenostylium recurvirostrum, 70

Hyophila involuta, 33; lusitanica, 15
Hypnella diversifolia, 224, 225*; key to the species, 224; leptorrhyncha, 224, 225*; pallescens, 224, 225*; pilifera, 224, 225*; punctata, 224*
Hypnum cupressiforme var. resupinatum, 33; diminutivum, *66*; pallescens "reptile expression," 33; palmatus, 37; pendulinum, 69; polypterum, 9; pungens, 26; reptans, 69; stenocarpum, 26; subpungens, 26; thelistegium, 68
Hypogymnia, 219+; atrofusca, 219; billardieri, 219; encausta, 222; imshaugii, 219; intestiniformis, 219+; mundata, 219; oroarctica, *222*; tubularis, 219
Hypogymniaceae, 222
hypogymnioid lichens, 219+
Hypotrachyna, 134

India, Rectolejeunea olivacea new to western Ghats, 195+; R. brittoniae new to India, 195+
insect dispersal, of Splachnum spores, 280
insects, role in distribution of the Splachnaceae, 279; taxa visiting Splachnum species, 282+
Interior Highlands, bryogeography of, 32; endemics in, 33
Isophyllaria attenuata, 292+
Isopterygiopsis muelleriana, 32
Isopterygium tenerum, 33, 60+
Isoviita, P. (review). H. Crum, Sphagnopsida. Sphagnaceae, 184
Iwatsuki, Z. A peculiar New Caledonian Sphagnum with rhizoids, 20
Iwatsukia, 11

Jaffueliobryum arsenei, *24*
Jamesoniella colorata, 293+
Jungermannia decurvifolia, *292*+; starkii, 155; torulosa, 186; tortuosa, 186; turneri, *161*

Keener, C. S. (review). V. H. Heywood & D. M. Moore (eds.), Current Concepts in Plant Taxonomy, 87
Kingiobryum paramicola, 9, 70
Koide, R. (review). V. Ahmadjian & S. Paracer, Symbiosis. An Introduction to Biological Associations, 318
Krause, Siberian moss collection, 276
Kurzia sylvatica, 32
Kymatocalyx, 155

Lasallia, 133; papulosa, 116
Lawrey, J. D. (review). K. A. Kershaw, Physiological Ecology of Lichens, 250
Lawrey, J. D. Biological role of lichen substances, 111
Lecanora, 110, 134; atra, 117; caesiorubella, 104; caesiorubella subsp. merrillii, 104; caesiorubella subsp. saximontana, 104; campestris, 136+; cinereofusca, 105; cinereofusca var. appalachensis, 137+; imshaugii, 136+; leptacina, 192+; pallida, 104; pallida group, 99; pelobotryon, 136; perplexa, 137+; pseudistera, 136; subfusca group, 104, 136+; subimmergens, 136
Lecidea, 100+; consentiens, 136; coroniformis, *260*; globifera, *261*; granulosa, 113; griseoatra, 192; lithophilaplana group, 104; luridella, *265*; nipponica, *266*; phaeops, 136; russellii, *271*; tuckermanii, nom. inval., 253, 274; vallesiaca, *274*
Lecidella bullata, 192
Lejeunea, 195; laetivirens, 33; lamaceriana subsp. gemminata, 33; olivacea, *195*; subolivacea, *195*
Lepidolaena menziesii, 292
Lepidopilidium laevisetum, 10; portoricensis, 10
Lepidozia chordulifera, 292; laevifolia, 294
Leprocaulon, 102
Leptobarbula, 227; berica, 228
Leptodontium, 19; albovaginatum, 15; subplanifolium, 15
Leptohymenium, 28; ehrenbergianum, *28*; patulum, *28*; tenue, 28, 29*
Leptoscyphus antarcticus, 292; expansus, 295; horizontalis, 293; patagonicus, 292+
Letharia, 134; columbiana, 117; vulpina, 117
Lethariella, 102; cladonioides, 107
Leucobryum albidum, 33, 60+
Leucodon, 9; julaceus, 60; peruvianus, 9; subgracilis, 9
Leucolejeunea clypeata, 33; unciloba, 33
Leucoloma mariei, 11
Lewinsky, J. & D. Griffin, III. Orthotrichum pallens Brid. new to Venezuela, 163
lichen chemistry in systematics, a critique of, 133
Lichen decipiens, *261*; intestiniformis, *222*
lichens, morphotypes, enzyme analysis in Umbilicaria muhlenbergii, 285; origin of chemical diversity in, 97; secondary products in systematic theory, 91; species-pairs in, 129+; substances in North America Psora species, 254
Lichina confinis, 117
Lichinaceae, 100
Lindigia, 230; aciculata, 230; debilis, 230
Lobaria amplissima, 134; laetivirens, 117; ornata, 134; pulmonaria, 117, 134; retigera, 134
Lopadium fuscum, 232+; puiggarii, 232+
Lophocolea heterophylla, 60+; palustris, 292+
Lophozia crispata, 294; leucorrhiza, 295

Macromitrium leprieuii, 11; ulophyllum, 10
Madotheca navicularis, 291
Magelospora, 134
Malaysia, Mitthyridium in, 49+
Marsupella sphacelata, 33
Mastigobryum fuegianum, 292
McIntosh, T. T. & J. R. Spence. Grimmia olympica Britt. (Grimmiaceae) is transferred to Brachydontium (Seligeriaceae), 200
Meesia craigieburnensis, *303*+
Melanelia calva, 296; glabratuloides, 296; subglabra, 296
Menegazzia, 219+
Metzgeria crassipilis, 32
mevalonic acid pathway, in lichen systematics, 91+
Mexico, Fissidens intermedius new to, 300; Gyroweisia monterreia in, 227; Leptohymenium tenue in, 28; Mittenothamnium in, 66; new species of Cygniella, Breedlovea, Hookeriopsis, Pyleisiadelpha in, 23+; Ozarkian elements in, 32
Micarea, 134
Microcampylopus, 152
Microthamnium mexicanum, *69*; pendulinum, *69*; pringlei, *66*
Mittenothamnium diminutivum, *66*; imbricatum, *66*; mexicanum, *69*; pendulinum, *69*; pringlei, *66*; reptans, 68*; squarrosulum, *68*; subthelistegum, *66*; volvatum var. mexicanum, *69*
Mitthyridium, 49+; adpressum, *52*; cardotii, 49+, 53*; constrictum, 49+, 55*; crassum, 49+, 56*; fasciculatum, 49+, 52*; fasciculatum var. rhizophyllum, *53*; flavum, 49, 50*; jungquilianum, 49+, 51*; key to species of Malaysia and adjacent regions, 49; luteum, 49+, 54*; manii, *49*; megamorphum, 49+; obtusifolium, 49+, 53*; parvifolium, *51*; perundulatum, *51*; repens, 49+, 51*; subluteum, 49+, 54*; undulatulum, *52*; undulatum, 49+, 52*; vriesii, *50*; wallisii, 49+, 55*
Moenkemeyera abrupta, *300*
Mohamed, D. *See* Reese, W. D., H. Mohamed & D. Mohamed, 49

MOHAMED, H. *See* REESE, W. D., H. MOHAMED & D. MOHAMED, 49
Molendoa obtusifolia, 9
Morinia, 13
MURRAY, B. M. Andreaea sinuosa, sp. nov. (Musci: Andreaeaceae), from Alaska, British Columbia and Scotland, 189
Mytilopsis albifrons, 11
Myurella sibirica, 33

Neckera complanata, 33
Neofuscelia, 299; adpicta, 296+; **atrobarbatica**, comb. nov., 297; brattii, 297; brunella, 297; delisei, 298; fissurina, 297; glabrans, 297; imitatrix, 298; loxodes, 115; **neodelisei**, comb. nov., 298; pulla, 296; squamans, 298; subimitatrix, 298; verrucella, 298; verruculifera, 114; waiporiensis, 298
Niagara River Gorge, 70
NISHIMURA, N. & H. ANDO. A revision of some Mittenothamnium species described from Mexico, 66
North America, Psora species new to, 253

Ochrolechia, 134; geminipara, 136; pacifica, 136; parella, 117; yasudae, 116
Octoblepharum, 10
Odontoschisma prostratum, 60
ORBAN, S. *See* REESE, W. D. & S. ORBAN, 213
Orthotrichum diaphanum, 33; gymnostomum, 165+; obtusifolium, 165+; pallens, 163*+; pallens var. crenulatum, 164; pallens var. johnsenae, 164; penicillatum, 163; pycnophyllum, 163+; rupestre, 163+

Pachyglossa exilis, 293; fissa, 293
Palamocladium leskeoides, 33
Pallavicinia, 292; pallidovirens, 292; pisicolor, 295
PALMER, M. W. Pattern in corticolous bryophyte communities of the North Carolina piedmont: do mosses see the forest or the trees?, 59
palmitic acid, 91
Pannaria ahlneri, 104
Paraparmelia, 297+; conranensis, 298
Parisia, 206+; ciliata, 208+, 211*; key to species of, 208; laevipila, 208+, 210*; neocaledonica, 206, 209*
Parmelia, 102+; atrobarbatica, *297*; borreri, 103; ceratophylla var. atrofusca, *222*; direagens, 107; hababiana, 107; neodelisei, *298*; perforata, 105; pulla group, 100; reddenda, 103; sorocheila, 129; stictica, 103; subrudecta, 103
Parmotrema dominicanum, 99; hypoleucinum, 127+; hypotropum, 127+; paramoreliense, 129; perforatum, 127+; perforatum complex, 127+; preperforatum, 127+
Pedinophyllopsis, 292; abditus, 294+
Peltigera aphthosa, 117, 134; canina, 117+; polydactyla, 117; praetextata, 117
Pertusaria, 100+; oculata, 107; pertusa, 115
Peru, Fissidens reesei new to, 300
Phaeophyscia, 102
phenolic acid units, biosynthetic origins of, 93
Philonotis elegantula, 9
phycobiont regulators, lichen secondary products as, 118
Physcia, 102; adscendens, 117
Physcomitrium, 46+; benoistii, *303*+; conicum, 302; laxum, *302*+
Pilopogon longirostratus, 10
Pilotrichella pentasticha, 11
Pilotrichidium callicostata, 10
Pinnatella pinniformis, 10
Pireella husnotiana, 10
Placopsis gelida, 117
Plagiochila abdita, *292*; acanthocaulis, 292; acanthophylla subsp. japonica var. ciligera, 33; angulata, 293; ansata, 295; ludoviciana, 33; molliuscula, 294; porelloides, 32; undata subsp. undata, 33
Plagiothecium, 46; lucidulum, 8
Platismatia, 102
Platygyrium repens, 60+
Pleistocene refugium, Interior Highlands as, 32
Pleurochaete squarrosa, 33
Pleurozium schreberi, 33
Pogonatum cuspidatum, 10
Pohlia filiformis, 33
Polytrichadelphus longisetus, 8
Polytrichum, 46, 144+; juniperinum, 144+; piliferum, 32
Porella platyphylla, 61
Pottia intermedia, 37
Prionodon densus, 11; fusco-lutescens, 10
Prionolobus turneri, *155*
Protoparmelia, 136
Protousnea, 102
Pseudephebe, 102
Pseudevernia, 220+; furfuracea, 105
Pseudocrossidium aureum, 72; replicatum, 10
Pseudocyphellaria anthrapsis, 117; flavicans, 117; intricata, 117
Pseudolepicolea, 292; quadrilaciniata, 295
Pseudoparmelia, 297; baltimorensis, 116, 128; caperata, 128
Pseudosymblepharis schimperiana, 9
Psora, 102, 253+; albilabra, *274*; **californica**, sp. nov., 254+, 257*; cerebriformis, 254+, 259*; concava, *253*+, *260*+; crenata, 254+, 260*; decipiens, 254+; 261*; elenkinii, *263*; globifera, 254+, 262*; **himalayana**, comb. nov., 254+, 262*; hypotheya, 265; icterica, 254+, 264*; lurida var. americana, *265*; luridella, 254+; mexicana, 253+; **montana**, sp. nov., 254+, 266*; nicolai, *253*+; nipponica, 253+, 267*; **nitida**, sp. nov., 254+, 268*; novomexicana, *253*+; **pacifica**, sp. nov., 254+, 268*; **pseudorussellii**, sp. nov., 254+, 269*; rubiformis, 253+, 271*; rufonigra f. terrena, 275; russellii, 254+, 272*; **tenuifolia**, sp. nov., 254+, 273*; **tuckermanii**, sp. nov., 253+, 273*; vallesiaca, 254+, 274*
Psorula rufonigra, *275*
Ptilidium pulcherrimum, 33
Ptychomitrium leibergii, 33
Punctelia, 102+; hypoleucites, 128; semansianum, 128
PURSELL, R. A. & D. W. VITAL. Distributional adumbrations of Fissidens in the neotropics, 300
PURSELL, R. A. (review). D. M. Greene, A Conspectus of the Mosses of Antarctica, South Georgia, the Falkland Islands and Southern South America, 318
PURSELL, R. A. Typification of Hedwig's species of Fissidens, 35
Pylaisiadelpha **sharpii**, sp. nov., 23*, 27; tenuirostris var. rhaphidostegioides, 27
Pyxine caesiopruinosa, 103

Racomitrium aciculare, 33; andreaeopsis, *277*+
Radula australis, 33; complanata, 61+; mollis, 33; obconica, 33, 61
RAEYMAEKERS, G. & J. M. GLIME (review). A. Schaepe, Veränderungen der Moosflora von Berlin (West), 318
Ramalina atlantica, 106, 126; crassa, 106, 126; cuspidata, 105+, 126; druidarum, 106; siliquosa, 105+, 126, 134; siliquosa group, 126; stenoclada, 106, 126; translucida, 107
rangiformic acid, 93

Rectolejeunea, 195; aloba, 195; arnellii, *198*; brittoniae, 195, 197*; **olivacea**, comb. nov., 195, 196*; subacuta, *195*
REDFEARN, P. L. Bryogeography of the Interior Highlands of North America: taxa of critical importance, 32
REESE, W. D. & S. ORBAN. Syrrhopodon incompletus Schwaegr. var. incompletus (Musci: Calymperaceae) disjunct between the Americas and Africa, 213
REESE, W. D., H. MOHAMED & D. MOHAMED. A synopsis of Mitthyridium (Musci: Calymperaceae) in Malaysia and adjacent regions, 49
Relicina, 102; fluorescens, 107
Rhabdogrimmia, 201
Rhabdoweisia crispata, 33
Rhaphidostichum elegans, 11; schwaneckeanum, 11; subluxurians, 11
Rhexophyllum, 13
rhizoids, in Sphagnum novo-caledoniae, 20
Rhizoplaca, 136
Rhynchostegiella, 230; attenuata, *230*; menadensis, 230; tenella, 230
Rhynchostegiopsis planifolia, 9
Rhytidiadelphus triquetrus, 33
Rhytidium rugosum, 33
Riccardia pallidovirens, 292+; papillosa, 292; prehensilis, 294
Riccia ozarkana, 33
ROBINSON, H. Notes on the bryogeography of Venezuela, 8
ROHRER, J. R. Leptohymenium tenue and Elmeriobryum guatemalense, sp. nov. in Mexico and Central America, 28
RUSHING, A. E. & Z. B. CAROTHERS. Comparative studies of spermatogenesis in the Bryopsida. III. Blepharoplast morphology in Thuidium delicatulum, 144

Schistidium andreaeopsis, 277*; holmenianum, *276+*
Schistochila alata, 293; laminigera, 292; splachnophylla, 292
Schistomitrium, 11
Schlotheimia fuscoviridis, 10
Schraderella, 26
Schraderobryum, 26; exiguum, 26; minutipes, 26; stenocarpum, *26*; subpungens, *26*; ulicinum, *26*; ulicinum var. exiguum, 26
Schwetschkeopsis fabronia, 33
Scopelophila ligulata, 33
Scotland, Andreaea sinnuosa in, 192
secondary chemical products, antimicrobial, allelopathic and antiherbivorous, 112; in hypogymnioid lichens, compared, 221
Seligeria, 200+; donniana, 33; pusilla, 33
Sematophyllum adnatum, 33, 60; marylandicum, 33; minutipes, 26; ulicinum, *26*
Sendtnera quadrilaciniata, *292*
Sharp, A. J., biography, 3
shikimic acid pathway, in lichen systematics, 91+
SMITH, C. W. Three foliicolous lichens new to the United States, 232
species-pairs, in lichens, 129
SPENCE, J. R. Bryum calobryoides, a new species from western North America, 215
SPENCE, J. R. *See* MCINTOSH, T. T. & J. R. SPENCE, 200
Sphaerothecium, 152; comosum, *152+*; phascoides, 152+, 153*+; reconditum, 152+, 153*; **subchlorophyllosum**, comb. nov., 152+, 154*
Sphagnum, 20, 144+; girgensohnii, 33; leratianum, *20*; novo-caledoniae, 21*, ecology of, 20; palustre, 144; perichaetiale, 20
Splachnaceae, role of insects in species distribution, 279+, in spore dispersal, 279+; substrate restriction in, 279
Splachnum, 9, 47, 282; ampullaceum, 279+; luteum, 279; rubrum, 279+; sphaericum, 279+; turbinatum, 9; vasculosum, 279+; weberbaueri, 9
Sporopodium phyllocharis, 232+
SRIVASTAVA, S. C. & A. AGARWAL. The genus Rectolejeunea in India, 195
Steere, W. C., dedicatory issue, Vol. 20 Evolutionary Biology, 252
STEERE, W. C. Trachyodontium, a new genus of the Pottiaceae (Musci) from Ecuador, 17
Stereocaulon, 102, 134; alpinum, 117; rivulorum, 107; vesuvianum, 117
Steyermarkiella anomalodictya, 11
Sticta sylvatica, 117
STOTLER, R. E. & B. CRANDALL-STOTLER. Recent literature on hepatics and anthocerotes—107, 73; —108, 168
Streptopogon erythrodontus, 10
Strigula elegans, 233
Sulcaria, 102
Sullivant, treatment of the Wilkes Expedition hepatics of, 291+
Symblepharis tenuis, 10
Symphyogyna crassifrons, *292*
Syrrhopodon, 49; adpressus, *52*; brevivaginans, *213*; crassus, *56*; crispatus, *15*, cryptocarpos, 9, 213; cuspidatus, *50*; fasciculatus var. rhizophyllus, *52*; flavus, *50*; gaudichaudii, 213; geheebii, *52*; glaucinus, *55*; gracilis, *52*; grandidens, *213*; humilis, *51*; incompletus var. incompletus, 213; jamesonii, 15; jungquilianus, *51*; leucoloma, *54*; louisiadum, *51*; luzonensis, *50*; nicobaricus, *50*; obtusifolius, *53*; obtusifolius var. neocaledonicus, *53*; ovatifolius, *50*; papuanus, *55*; perundulatus, *51*; repens, *51*; subluteus, *55*; subobtusifolius, *54*; subundulatus, *53*; texanus, 33; undulatulus, *51*; vriesii, *50*; wallisii, *56*
systematics, lichen secondary products in, 91

Tapellaria epiphylla, 233
Telaranea blepharostoma, 294; oligophylla, 294; pseudozooposis, 293
Temnoma quadripartitum, 293+
Tephromela, 136
Tetraphis, 47; pellucida, 33, 149
Tetraplodon mnioides, 11, 279+
Thamnolia subuliformis, 105; vermicularis, 105
Thelia asprella, 61
Thelotremataceae, 100
THIERS, B. M. (review). S. Piippo, A Monograph of the Genera Lepidolejeunea and Luteolejeunea (Lejeuneaceae, Hepaticae), 248
THIERS, B. M. (review). Y. Kuwahara, The Metzgeriaceae of the Neotropics, 247
Thuidium, 47, 144+; blepharoplast morphology in, 144+, 146*, 147*, 148*; delicatulum, 60+; minutulum, 33
Thyridium adpressum, *52*; cardotii, *55*; cuspidatum, *50*; flavum var. ovatifolium, *50*; geheebii, *52*; louisiadum, *51*; luteum, *55*; luzonensis, *50*; megamorphum, *57*; nicobaricum, *50*; parvifolium, *51*; perundulatum, *51*; samoanum, *54*; undulatulum, *52*; vriesii, *50*
Thysanomitrion comosum, *152*; phascoides, *152+*
TIMDAL, E. A revision of Psora (Lecideaceae) in North America, 253
Timmia megapolitana, 33
Timmiella barbuloides, 149
Toninia tristis, 268
Tortella humilis, 60

Tortula arcuata, *15*; fragilis, 33; pagorum, 33; papillosa, 33; rhizophylla, 71; ruralis, 33
Trachyodontium zanderi, gen. et sp. nov., 17*
TRAVERSE, A. (review). A. Gupta & R. Udar, Palyno-Taxonomy of Selected Indian Liverworts, 249
Trebouxia, 134
Trematodon longicollis, 33
Triandrophyllum subtrifidum, 294
Trichostomopsis umbrosa, *70*+
Trichostomum amblystegium, *15*; arcuatum, *15*; campylocarpum, *15*; campylopyxis, *15*
Trigonanthus divaricatus, 155
Tuerckheimia angustifolia, 33
Tylimanthus flavicans, 293; urvilleanus, 294

Ulota hutchinsiae, 33
Umbilicaria mammulata, 139, 289; muhlenbergii, 285+, 287*; vellea, 285+
United States, three lichens new to, 232
Usnea antarctica, 107; florida, 134; strigosa, 107+; subfloridana, 117, 139
usnic acid, antibiotic properties of, 112

Venezuela, African floristic elements in, 11; bryogeography of, 8; Orthotrichum pallens new to, 163
Venturiella sinensis var. angusti-annulata, 33
VITAL, D. M. *See* PURSELL, R. A., 300

Weber, W. A., Robert L. Stearns Award given to, 252
Weissia condensa, 227; jamaicensis, 33; sharpii, 33
West Indies, endemism in, 11
WHITMORE, A. (review). S. Jovet-Ast, Les Riccia de la Region Mediterranee, 317
Wilkes U.S. Exploring Expedition, hepatic collections of, 291
WILSON, P. Homalothecium arenarium (Lesq.) Lawt. illustrated, 328
wind dispersal, of Splachnum spores, 280
WYATT, R. *See* CAMERON, R. G. & R. WYATT, 279

Xanthoparmelia arida, 100; arseneana, 100; barbatica, 100; cumberlandia, 100, 116; dissensa, 100; lineola, 100; novomexicana, 100; psoromifera, 100; taractica, 100; tucsonensis, 100
Xanthoria parietina, 115
Xanthospora, 102
xerothermal period, dispersal into the Interior Highlands, 33

ZANDER, R. H. & F. J. HERMANN. Gyroweisia monterreia, a new moss species from Mexico, 227
ZANDER, R. H. Notes on Bryoerythrophyllum (Musci), 13
Zygodon apiculatus, 33

THE BRYOLOGIST

Quarterly Journal of the American Bryological and Lichenological Society

Editor
DOMINICK V. BASILE

Volume 89
1986

Table of Contents

AHMADJIAN, V. Review 185
ALLEN, B. H. A Review of the Genus *Parisia* (Musci: Dicranaceae) 206
ALLEN, B. H. The Taxonomic Status of *Hypnella punctata* 224
ALLEN, B. H. Reviews 185, 248
BRODO, I. M. Interpreting Chemical Variation in Lichens for Systematic Purposes 132
BRODO, I. M. & W. L. CULBERSON. *Haematomma pustulatum,* sp. nov. (Ascomycotina, Haematommataceae): A Common, Widespread, Sterile Lichen of Eastern North America 203
BROWN, R. C. & Z. B. CAROTHERS. Comparative Studies of Spermatogenesis in the Bryopsida. II. Blepharoplast Morphology in *Archidium tenerrimum* Mitt. 42
BUCK, W. R. Reviews 249, 250
CAMERON, R. G. & R. WYATT. Substrate Restriction in Entomophilous Splachnaceae: Role of Spore Dispersal 279
CROSBY, M. R. & C. R. BAUER. Recent Literature on Mosses—135 177
CRUM, H. Taxonomic and Nomenclatural Addenda to the Mexican Moss Flora 23
CULBERSON, C. F. Biogenetic Relationships of the Lichen Substances in the Framework of Systematics 91
CULBERSON, W. L. Natural-Product Chemistry in the Systematics of the Lichen Fungi. Introduction 89
CULBERSON, W. L. Chemistry and Sibling Speciation in the Lichen-Forming Fungi: Ecological and Biological Considerations 123
CULBERSON, W. C. Reviews 86, 87
DELGADILLO M., C. Aaron J. Sharp: A Life-time of Botanical Experience 3
ECKEL, P. M. *Didymodon australasiae* var. *umbrosus* New to Eastern North America 70
EGAN, R. S. Correlations and Non-Correlations of Chemical Variation Patterns with Lichen Morphology and Geography 99
EGAN, R. S. Recent Literaure on Lichens—126, 127, 128 ..... 80, 234, 310
ESSLINGER, T. L. Notes on the Chemistry and Distribution of Selected Parmeliaceae (Lichens) from the Southern Hemisphere 296
FAHSELT, D. Multiple Enzyme Forms of Morphotypes in a Population of *Cladonia cristatella* Tuck. 139
FIFE, A. J. Taxonomic and Nomenclatural Observations on the Funariaceae. 4. A Review of *Entosthodon laxus* with Incidental Notes on *E. obtusifolius* 302
FRAHM, J.-P. A Review of *Sphaerothecium* Hampe 152
FRISVOLL, A. A. The Status of *Schistidium andreaeopsis,* with Notes on the Krause Collection from the Chuck Peninsula 276
GOWARD, T. *Brodoa,* a New Lichen Genus in the Parmeliaceae 219
GRIFFIN, III, D. *Rhynchostegiella attenuata* Batr. (Brachytheciaceae): A Synonym of *Lindigia aciculata* (Tayl.) Hampe (Meteoriaceae) 230
HAGEMAN, C. & D. FAHSELT. A Comparison of Isozyme Patterns of Morphological Variants in the Lichen *Umbilicaria muhlenbergii* (Ach.) Tuck. 285
HASSEL DE MENENDEZ, G. G. The Discovery of Sullivant's Wilkes U.S. Exploring Expedition Hepaticae. The "Fuegian" Specimens 291
HEDDERSON, T. A. A Naturally Occurring Moss Hybrid Between *Orthotrichum gymnostomum* and *O. obtusifolium* 165
HONG, W. S. The Family Cephaloziellaceae in North America West of the Hundredth Meridian 155
ISOVIITA, P. Review 184
IWATSUKI, Z. A Peculiar New Caledonian *Sphagnum* with Rhizoids 20
KEENER, C. S. Review 87
KOIDE, R. Review 319
LAWREY, J. D. Biological Role of Lichen Substances 111
LAWREY, J. D. Review 250
LEWINSKY, J. & D. GRIFFIN, III. *Orthotrichum pallens* Brid. New to Venezuela 163
MCINTOSH, T. T. & J. R. SPENCE. *Grimmia olympica* Britt. (Grimmiaceae) is Transferred to *Brachydontium* (Seligeriaceae) 200

MURRAY, B. M. *Andrea sinuosa,* sp. nov. (Musci: Andreaeaceae), from Alaska, British Columbia and Scotland 189
NISHIMURA, N. & H. ANDO. A Revision of Some *Mittenothamnium* Species Described from Mexico 66
PALMER, M. W. Pattern in Corticolous Bryophyte Communities of the North Carolina Piedmont: Do Mosses See the Forest or the Trees? 59
PURSELL, R. A. Typification of Hedwig's Species of *Fissidens* 35
PURSELL, R. A. & D. M. VITAL. Distributional Adumbrations of *Fissidens* in the Neotropics 300
PURSELL, R. A. Review 318
RAEYMAEKERS, G. & J. M. GLIME. Review 318
REDFEARN, JR., P. L. Bryogeography of the Interior Highlands of North America: Taxa of Critical Importance 32
REESE, W. D., H. MOHAMED & A. D. MOHAMED. A Synopsis of *Mitthyridium* (Musci: Calymperaceae) in Malaysia and Adjacent Regions 49
REESE, W. D. & S. ORBAN. *Syrrhopodon incompletus* Schwaegr. var. *incompletus* (Musci: Calymperaceae) Disjunct Between the Americas and Africa 213
ROBINSON, H. Notes on the Bryogeography of Venezuela 8
ROHRER, J. R. *Leptohymenium tenue* and *Elmeriobryum guatemalense,* sp. nov. in Mexico and Central America 28
RUSHING, A. E. & Z. B. CAROTHERS. Comparative Studies of Spermatogenesis in the Bryopsida. III. Blepharoplast Morphology in *Thuidium delicatulum* 144
SMITH, C. W. Three Foliicolous Lichens New to the United States 232
SPENCE, J. R. *Bryum calobryoides,* a New Species from Western North America 215
SRIVASTAVA, S. C. & A. AGARWAL. The Genus *Rectolejeunea* in India 195
STEERE, W. C. *Trachyodontium,* a New Genus of the Pottiaceae (Musci) from Ecuador 17
STOTLER, R. E. & B. CRANDALL-STOTLER. Recent Literature on Hepatics and Anthocerotes—107, 108 ..... 73, 168
THIERS, B. M. Reviews 247, 248
TIMDAL, E. A Revision of *Psora* (Lecideaceae) in North America 253
TRAVERSE, A. Review 249
WHITMORE, A. Review 317
ZANDER, R. H. Notes on *Bryoerythrophyllum* (Musci) 13
ZANDER, R. H. & F. J. HERMANN. *Gyroweisia monterreia,* a New Moss Species from Mexico 227

Actual dates of issue of Volume 89:
Number 1: 27 June 1986
Number 2: 8 October 1986
Number 3: 29 January 1987
Number 4: 20 May 1987